

**RESQPOD® ITD BIBLIOGRAPHY**SELECTED STUDIES THAT INCLUDED USE OF AN IMPEDANCE THRESHOLD DEVICE (ITD)  
DURING THE PERFORMANCE OF MANUAL OR AUTOMATED CPR**Clinical (Human) Studies**

1. Aufderheide TP, Pirrallo RG, Provo TA, Lurie KG. Clinical evaluation of an inspiratory impedance threshold device during standard cardiopulmonary resuscitation in patients with out of hospital cardiac arrest. *Crit Care Med.* 2005;33:734-740.
2. Pirrallo RG, Aufderheide TP, Provo TA, Lurie KG. Effect of an inspiratory impedance threshold device on hemodynamics during conventional manual cardiopulmonary resuscitation. *Resuscitation.* 2005;66:13-20.
3. Thayne R, Thomas DC, Neville JD, Van Dellen A. Use of an impedance threshold device improves short-term outcomes following out-of-hospital cardiac arrest. *Resuscitation.* 2005;67(1):103-108.
4. Aufderheide TP, Alexander C, Lick C, et al. From laboratory science to six emergency medical services systems: new understanding of the physiology of cardiopulmonary resuscitation increases survival rates after cardiac arrest. *Crit Care Med.* 2008;36:S397-S404.
5. Cabrini L, Beccaria P, Landoni G, et al. Impact of impedance threshold devices on cardiopulmonary resuscitation: a systematic review and meta-analysis of randomized controlled studies. *Crit Care Med.* 2008;36(5):1625-1632.
6. Lurie K, Schnettler P, Steinkamp J, et al. Level One Cardiac Arrest Centers are clinically and cost effective. *Circulation.* 2008;118:S1486.
7. Aufderheide TP, Lurie KG, Yannopoulos D, et al. Ventilation rate and use of the impedance threshold device are correlated with hemodynamics during CPR in humans. *Circulation.* 2009;120:S-669.
8. Aufderheide TP, Yannopoulos D, Lick CJ, et al. Implementing the 2005 American Heart Association guidelines improves outcomes after out-of-hospital cardiac arrest. *Heart Rhythm.* 2010;7(10):1357-1362.
9. Dailey M, Politis J, Provo TA. Implementation of the AHA guidelines with a systems-based approach improves survival to hospital discharge following cardiac arrest. *Circulation.* 2010;122:A51.
10. Hinchey PR, Myers JB, Lewis R, et al for the Capitol County Research Consortium. Improved out-of-hospital cardiac arrest survival after the sequential implementation of the 2005 AHA guidelines for compressions, ventilations and induced hypothermia: the Wake County experience. *Ann Emerg Med.* 2010;56(4):358-361.
11. Saussy JM, Elder JE, Flores CA, Miller AL. Optimization of cardiopulmonary resuscitation with an impedance threshold device, automated compression cardiopulmonary resuscitation and post-resuscitation in-the-field hypothermia improves short-term outcomes following cardiac arrest. *Circulation.* 2010;122:A256.
12. Thigpen K, Davis SP, Basol R, et al. Implementing the 2005 American Heart Association guidelines, including use of an impedance threshold device, improves hospital discharge rates after in-hospital cardiac arrest. *Respir Care.* 2010;55(8):1014-1019.
13. Thigpen K, Simmons L, James Z, Neely C. Implementation of the 2005 American Heart Association guidelines improves in-hospital cardiac arrest survival rates in a community hospital: a 5-year case series. *Circulation.* 2010;122:A48.
14. Lick CJ, Aufderheide TP, Niskanen RA, et al. Take Heart™ America: a comprehensive, community-wide, systems-based approach to the treatment of cardiac arrest. *Crit Care Med.* 2011;39(1):26-33.

15. Aufderheide TP, Nichol G, Rea TD, et al for the ROC investigators. A trial of an impedance threshold device in out-of-hospital cardiac arrest. *NEJM*. 2011;365:798-806.
16. Thigpen K, Simmons L, James Z, Neely C. Breaking the 30% survival rate window: impact of the 2005 and 2010 American Heart Association guidelines on in-hospital cardiac arrest survival with favorable neurological function. *Circulation*. 2013;128:A164.
17. Schultz JC, Caldwell E, Sarraf M, et al. Prolonged adult cardiac arrest with enhanced cardiopulmonary resuscitation and sodium nitroprusside: a case series. *Circulation*. 2013;128:A248.
18. Satterlee PA, Boland LL, Johnson PJ, et al. Implementation of a mechanical chest compression device as standard equipment in a large metropolitan ambulance service. *J Emerg Med*. 2013;45(4):562-569.
19. Chase D, Salvucci A, Marino R, et al. Effect of airway management and impedance threshold device on circulation, survival and neurological outcome in adult out-of-hospital cardiac arrest. *Open J Emerg Med*. 2014;2:12-18.
20. Driver BE, Debaty G, Plummer DW, Smith SW. Use of esmolol after failure of standard cardiopulmonary resuscitation to treat patients with refractory ventricular fibrillation. *Resuscitation*. 2014;85:1337-1341.
21. Escott MEA, Jenks SP, Traynor KM, et al. External cardiac bypass? A case series: hemodynamics of LUCAS device plus an ITD in cardiac arrest. *Prehosp Emerg Care*. 2014;18(1):156.
22. Yannopoulos D, Aufderheide TP, Abella BS, et al. Quality of CPR: an important effect modifier in cardiac arrest clinical outcomes and intervention effectiveness trials. *Resuscitation*. 2015;94:106-113.
23. Idris AH, Guffey D, Pepe PE, et al for the ROC Investigators. Chest compression rates and survival following out-of-hospital cardiac arrest. *Crit Care Med*. 2015;43(4):840-846.
24. Yannopoulos D, Bartos JA, Martin C, et al. Minnesota Resuscitation Consortium's advanced perfusion and reperfusion cardiac life support strategy for out-of-hospital refractory ventricular fibrillation. *J Am Heart Assoc*. 2016;5(6):e003732.
25. Hopkins C, Burk C, Moser S, et al. Implementation of pit crew approach and cardiopulmonary resuscitation metrics for out-of-hospital cardiac arrest improves patient survival and neurological outcomes. *J Am Heart Assoc*. 2016;5(1):1-10.
26. Sporer K, Jacobs M, Derevin L, et al. Continuous quality improvement efforts increase survival with favorable neurological outcome after out-of-hospital cardiac arrest. *Prehosp Emerg Care*. 2016;1-6.
27. Sugiyama A, Duval S, Nakamura Y, et al. Impedance threshold device combined with high-quality cardiopulmonary resuscitation improves survival with favorable neurological function after witnessed out-of-hospital cardiac arrest. *Circ J*. 2016;80(10):2124-2132.
28. Adabag S, Hodgson L, Garcia S, et al. Outcomes of sudden cardiac arrest in a state-wide integrated resuscitation program: results from the Minnesota Resuscitation Consortium. *Resuscitation*. 2017;110:95-100.
29. Ma L, Xiong B. Effect of impedance threshold devices on cardiopulmonary resuscitation: an updated meta-analysis. *Crit Care Med*. 2018;46(1):118.
30. Duval S, Pepe PE, Aufderheide TP, et al. Optimal combination of compression rate and depth during cardiopulmonary resuscitation for functionally favorable survival. *JAMA Cardiol*. 2019;4(9):900-908.
31. Gonzales L, Oyler B, Hayes JL, et al. Out-of-hospital cardiac arrest outcomes with "pit-crew" resuscitation and scripted initiation of mechanical CPR. *Am J Emerg Med*. 2019;37:913-920.
32. Pepe PE, Schepke KA, Antevy PM, et al. Confirming the clinical safety and feasibility of a bundled methodology to improve cardiopulmonary resuscitation involving a head-up/torso-up chest compression technique. *Crit Care Med*. 2019;47:449-455.

## Animal Studies

1. Lurie KG, Mulligan KA, McKnite S, et al. Optimizing standard cardiopulmonary resuscitation with an inspiratory impedance threshold valve. *Chest*. 1998;113(4):1084-1090.
2. Lurie KG, Ingbar D, Sweeney M, et al. Potential role of the vocal cords during cardiopulmonary resuscitation. *Circulation*. 1998;98(17):I-478.
3. Lurie KG, Voelckel WG, Zielinski T, et al. Improving standard cardiopulmonary resuscitation with an inspiratory impedance threshold valve in a porcine model of cardiac arrest. *Anesth Analg*. 2001;93(3):649-655.
4. Langhelle A, Stromme T, Sunde K, et al. Inspiratory impedance threshold valve during CPR. *Resuscitation*. 2002;52:39-48.
5. Lurie KG, Zielinski T, McKnite S, et al. Use of an inspiratory impedance valve improves neurologically intact survival in a porcine model of ventricular fibrillation. *Circulation*. 2002;105(1):124-129.
6. Lurie KG, Barnes TA, Zielinski TM, McKnite SH. Evaluation of a prototypic inspiratory impedance threshold valve designed to enhance the efficiency of cardiopulmonary resuscitation. *Respir Care*. 2003;48(1):52-57.
7. Dorph E, Wik L, Stromme TA, et al. Oxygen delivery and return of spontaneous circulation with ventilation:compression ratio 2:30 versus chest compressions only CPR in pigs. *Resuscitation*. 2004;60:309-318.
8. Yannopoulos D, Sigurdsson G, McKnite S, et al. Reducing ventilation frequency combined with an inspiratory impedance device improves CPR efficiency in a swine model of cardiac arrest. *Resuscitation*. 2004;(61):75-82.
9. Yannopoulos D, Aufderheide TP, Gabrielli A, et al. Clinical and hemodynamic comparison of 15:2 and 30:2 compression-to-ventilation ratios for cardiopulmonary resuscitation. *Crit Care Med*. 2006;34(5):1444-1449.
10. Alexander C, Yannopoulos D, Aufderheide T, et al. Dual mechanism of blood flow augmentation to the brain using an impedance threshold device in a pediatric model of cardiac arrest. *Circulation*. 2007;116(16):II-433.
11. Aufderheide TP, Alexander C, Lick C, et al. From laboratory science to six emergency medical services systems: new understanding of the physiology of cardiopulmonary resuscitation increases survival rates after cardiac arrest. *Crit Care Med*. 2008;36:S397-S404.
12. Lurie KG, Yannopoulos D, McKnite SH, et al. Comparison of a 10-breaths-per-minute versus a 2-breaths-per-minute strategy during cardiopulmonary resuscitation in a porcine model of cardiac arrest. *Respir Care*. 2008;52(7):862-870.
13. Matsuura T, McKnite S, Metzger A, et al. An impedance threshold device combined with an automated active compression decompression CPR device (LUCAS) improves the chances for survival in pigs in cardiac arrest. *Circulation*. 2008;118:S1449-1450.
14. Buckley GJ, Shih A, Garcia-Pereira FL, Bandt C. The effect of using an impedance threshold device on hemodynamic parameters during cardiopulmonary resuscitation in dogs. *J Vet Emerg Crit Care*. 2012;22(4):435-440.
15. Burnett A, Salzman JG, Segal N, et al. Potential negative effects of epinephrine on carotid blood flow and ETCO<sub>2</sub> during active compression-decompression CPR utilizing an impedance threshold device. *Resuscitation*. 2012;83(8):1021-1024.
16. Debaty G, Segal N, Matsuura T, et al. Hemodynamic improvement of a LUCAS 2 automated device by addition of an impedance threshold device in a pig model of cardiac arrest. *Resuscitation*. 2014;85(12):1704-1707.

17. Debaty G, Shin SD, Metzger A, et al. Tilting for perfusion: head-up position during cardiopulmonary resuscitation improves brain flow in a porcine model of cardiac arrest. *Resuscitation*. 2015;87:38-43.
18. Segal N, Metzger AK, Moore JC, et al. Correlation of end tidal carbon dioxide, amplitude spectrum area, and coronary perfusion pressure in a porcine model of cardiac arrest. *Physiol Rep*. 2017;5(17):e13401.

## Human Cadaver Studies

1. Holley J, Moore JC, Jacobs M, et al. Supraglottic airway devices variably develop negative intrathoracic pressures: a prospective cross-over study of cardiopulmonary resuscitation in human cadavers. *Resuscitation*. 2020;148:32-38.

## Manikin/Bench Studies

1. Lurie KG, Barnes TA, Zielinski TM, McKnite SH. Evaluation of a prototypic inspiratory impedance threshold valve designed to enhance the efficiency of cardiopulmonary resuscitation. *Respir Care*. 2003;48(1):52-57.
2. Babbs CF. Effects of an impedance threshold valve upon hemodynamics in standard CPR: studies in a refined computational mode. *Resuscitation*. 2005;66(3):335-345.
3. Genzwuerker HV, Gernoth C, Hinkelbein J, et al. Influence of an impedance threshold valve on ventilation with supraglottic airway devices during cardiopulmonary resuscitation in a manikin. *Resuscitation*. 2010;81:1010-1013.

## Review Articles

1. Schultz JC, Yannopoulos D. Is intrathoracic pressure regulation at the threshold of new resuscitation science? *Crit Care Med*. 2012;40(3):1008-1009.
2. Debaty D, Metzger A, Lurie K. Evaluation of Zoll Medical's ResQCPR System for cardiopulmonary resuscitation. *Expert Rev Med Devices*. 2015;12(5):505-516.
3. Lurie KG, Nemergut EC, Yannopoulos D, Sweeney M. The physiology of cardiopulmonary resuscitation. *Anesth Analg*. 2016;122(3):767-783.
4. Riess, ML. New developments in cardiac arrest management. *Adv Anesth*. 2016;34:29-46.
5. Pepe PE, Aufderheide TP. EBM vs EBM: combining evidence-based and experienced-based medicine in resuscitation research. *Curr Opin Crit Care*. 2017;23(3):199-203.
6. Segal N, Youngquist S, Lurie K. Ideal (i)CPR: looking beyond shadows in a cave. *Resuscitation*. 2017;121:8183.

Pre-clinical results are not necessarily representative of clinical outcomes. The studies listed in this document are not intended to imply specific efficacy or safety claims not yet approved by the US FDA. The generally cleared indication for the ResQPOD ITD available for sale in the United States (US) is for a temporary increase in blood circulation during emergency care, hospital, clinic, and home use.